

Tropicoperdix charltoni, Blyth, J. As. Soc. Beng. xxviii. p. 415 (1859).

"*Peloperdix charltoni*, Blyth," Gray, List B. pt. v., Gall. p. 58 (1867).

Arborophila charltoni, Hume, Str. F. ii. p. 450 (1874).

"*Arboricola pyrrhogaster*, Reichenb.," Blyth, Cat. Mamm. & B. Burma, p. 151 (1875).

Hab. The Malay Peninsula from Penang southwards, Sumatra, and N.E. Borneo. Mr. Hume states that it has also been sent from Bangkok, in Siam, and that it is reported to have been obtained in the Southern Tenasserim Hills.

In the females the blackish-brown barring on the lower half of the chest is more broken up and less sharply defined than in the males.

XXXIV.—*Further Additions to the Avifauna of Tsu-sima, in the Japanese Empire.* By HENRY SEEBOHM.

MR. HOLST has sent me a third collection of birds from the island of Tsu-sima, in the Straits of Corea, which enables me to add five more species to the list of birds inhabiting the island, as given above, pp. 87, 248. As before, the numbers refer to my 'Birds of the Japanese Empire.'

132. PICUS RICHARDSI.

In addition to the two males of this fine Woodpecker sent in the previously-mentioned package, Mr. Holst has now sent two females shot on Tsu-sima, the one on the 14th and the other on the 30th of November. The extent of white at the tip of the outer web of the third primary is one inch in one of them and an inch and an eighth in the other. There can now be no doubt of the identity of *Picus kalinowskii* from Corea with this species.

170. SCOPS JAPONICUS.

Mr. Holst has sent an example of the Japanese race of the Scops Owl shot on Tsu-sima on the 27th of October. It is

rather grey, and the wing measures $5\frac{3}{4}$ inches from the carpal joint.

190. ACCIPITER PALUMBARIUS.

Mr. Holst has sent a fine female Goshawk, shot on Tsu-sima on the 26th of October.

193. PHALACROCORAX CARBO.

Mr. Holst has sent the skin of a Common Cormorant, which was shot on Tsu-sima on the 6th of November. It has fourteen tail-feathers, and the scapulars and wing-coverts are bronzy brown, margined with black. The Shags have only twelve rectrices.

246. FULIGULA CLANGULA.

Mr. Holst has sent an example of the Golden-eye, shot on Tsu-sima in November.

361. CREX FUSCA ERYTHROTHORAX.

Mr. Holst has sent an example of the Siberian race of the Ruddy Crake shot on Tsu-sima. It measures $4\frac{1}{2}$ inches in length of wing from the carpal joint.

XXXV.—*On Birds collected or observed in the Vicinity of Foochow and Swatow in South-eastern China.* By JOHN D. DE LA TOUCHE.—Part I. (Communicated by H. H. SLATER.)

THE province of Fukien is one of the smallest and, I believe, one of the wildest and least populated provinces of China. During the three and a half years which I spent at Foochow, my collecting-grounds were confined to the strip of country extending from the sea-coast to about 130 miles up the River Min. A brief account of the country may be of interest in connection with the list of birds obtained in the district.

From Ching-feng-ling*, the furthest point reached by

* I do not know the altitude of the Ching-feng Mountains, but I believe that the country stands at least 3000 feet above the sea-level, as for 15 miles before getting there the road goes on gradually rising till Ching-feng-ling is reached.